arXiv:1703.04508v1 [math.CO] 13 Mar 2017

PATTERN RECOGNITION ON ORIENTED MATROIDS: DECOMPOSITIONS OF TOPES, AND DEHN–SOMMERVILLE TYPE RELATIONS

ANDREY O. MATVEEV

Abstract. If $\mathrm{V}(\boldsymbol{R})$ is the vertex set of a symmetric cycle $\boldsymbol{R}$ in the tope graph of a simple oriented matroid $\mathcal{M}$, then for any tope T of $\mathcal{M}$ there exists a unique inclusion-minimal subset $\boldsymbol{Q}(T,\boldsymbol{R})$ of $\mathrm{V}(\boldsymbol{R})$ such that T is the sum of the topes of $\boldsymbol{Q}(T,\boldsymbol{R})$.

If $|\boldsymbol{Q}(T,\boldsymbol{R})|\ge 5$, then the decomposition $\boldsymbol{Q}(T,\boldsymbol{R})$ of the tope T with respect to the symmetric cycle $\boldsymbol{R}$ satisfies certain Dehn–Sommerville type relations.

1. Introduction

Let $\mathcal{A}:=(E_t,\mathcal{L})=(E_t,\mathcal{T})$ be a *simple* (i.e., with no *loops*, *parallel elements* or *antiparallel elements*) *acyclic* oriented matroid on the ground set $E_t:=\{1,2,\dots,t\}$, with set of covectors $\mathcal{L}$ and with set of topes $\mathcal{T}\subseteq\{1,-1\}^{E_t}$, of rank $r\ge 3$. Let $\boldsymbol{R}$ be a *symmetric* $2t$*-cycle*, with its vertex set $\mathrm{V}(\boldsymbol{R})=-\mathrm{V}(\boldsymbol{R})$, in the tope graph $\mathcal{T}(\mathcal{L}(\mathcal{A}))$ of $\mathcal{A}$ such that $(1,1,\dots,1)=:\mathrm{T}^{(+)}\notin\mathrm{V}(\boldsymbol{R})$.

Recall that the vertex set $\mathrm{V}(\boldsymbol{R})$ of the cycle $\boldsymbol{R}$ is the set of topes of a *rank* 2 *oriented matroid* denoted $\mathcal{A}_{\boldsymbol{R}}$, see e.g. [3, Rem. 1.7], and thus by [1, Example 7.1.7] we can consider the representation of $\mathcal{A}_{\boldsymbol{R}}$ by a certain central line arrangement

$$\{\mathbf{x}\in\mathbb{R}^2\colon\langle\boldsymbol{a}_e,\mathbf{x}\rangle=0,\ e\in E_t\}\tag{1.1}$$

in the plane $\mathbb{R}^2$, with the set of corresponding normal vectors

$$\boldsymbol{A}:=\{\boldsymbol{a}_e\colon e\in E_t\}\,,\tag{1.2}$$

and with the standard scalar product $\langle\cdot,\cdot\rangle$.

For any tope T of the oriented matroid $\mathcal{A}$ there exists a *unique inclusion-minimal subset* $\boldsymbol{Q}(T,\boldsymbol{R})\subset\mathrm{V}(\boldsymbol{R})$ such that

$$\sum_{Q\in\boldsymbol{Q}(T,\boldsymbol{R})}Q=T\,,$$

see [3, Sect. 11.1].

If the tope graph $\mathcal{T}(\mathcal{L}(\mathcal{A}))$ is the *hypercube graph* $\boldsymbol{H}(t,2)$ on 2^t vertices, then for any *odd* integer j, $1\le j\le t$, by [3, Th. 13.6] there are precisely $2\binom{t}{j}$ vertices T of $\boldsymbol{H}(t,2)$ such that $|\boldsymbol{Q}(T,\boldsymbol{R})|=j$.

Let $\mathbf{max}^+ \mathrm{V}(\boldsymbol{R})$ denote the subset of topes of $\mathrm{V}(\boldsymbol{R})$ with *inclusion-maximal positive parts*. For the *positive tope* $\mathrm{T}^{(+)}$ of $\mathcal{A}$, by [3, Prop. 1.12(ii)] we have

$$\boldsymbol{Q}(\mathrm{T}^{(+)}, \boldsymbol{R}) = \mathbf{max}^+ \mathrm{V}(\boldsymbol{R}) \, .$$

If

$$|\boldsymbol{Q}(\mathrm{T}^{(+)}, \boldsymbol{R})| \geq 5 \, ,$$

then for the set of vectors (1.2) we have

$$|\{\boldsymbol{a} \in \boldsymbol{A} \colon \boldsymbol{a} \in \mathbf{C}_{>}\}| \geq 2 \, , \tag{1.3}$$

for any *open half-space* $\mathbf{C}_{>} \subset \mathbb{R}^2$ bounded by a 1-dimensional subspace of $\mathbb{R}^2$; see e.g. [2, Proof of Prop. 2.33].

Denote by ν_j the number of *feasible* subsystems, of cardinality j, of the *infeasible system* of *homogeneous strict linear inequalities*

$$\left\{ \langle \boldsymbol{a}_e, \mathbf{x} \rangle > 0 \colon \mathbf{x} \in \mathbb{R}^2, \ \boldsymbol{a}_e \in \boldsymbol{A} \right\} \tag{1.4}$$

associated with the arrangement (1.1). Since the condition (1.3) holds, the quantities ν_j satisfy the relations (where x is a formal variable)

$$\begin{cases} \nu_j = \binom{t}{j}, \text{ if } 0 \leq j \leq 2 \, , \\ \nu_{t-1} = \nu_t = 0 \, , \\ \sum_{j=3}^{t} \left(\binom{t}{j} - \nu_j \right) (\mathrm{x}-1)^{t-j} = - \sum_{j=3}^{t} (-1)^j \left(\binom{t}{j} - \nu_j \right) \mathrm{x}^{t-j} \end{cases} \tag{1.5}$$

called the *Dehn–Sommerville equations for the feasible subsystems of the system* (1.4); see e.g. [2, Prop. 3.53].

In particular,

- if $t = 5$, then
$$\nu_3 = \tbinom{t}{2} - t \, ;$$
- if $t = 6$, then
$$\nu_3 = \tbinom{t}{3} - 2t + 4 \, ,$$
$$\nu_4 = \tbinom{t}{2} - 3t + 6 \, ;$$
- if $t = 7$, then
$$\nu_4 = 2\nu_3 - 2\tbinom{t}{4} + \tbinom{t}{3} \, ,$$
$$\nu_5 = \nu_3 - \tbinom{t}{4} + \tbinom{t}{2} - t \, .$$

For any j, $3 \leq j \leq t-2$, by [2, Cor. 3.55(i)] we have

$$\tbinom{t}{j} - \nu_j = - \sum_{i=3}^{j} (-1)^i \tbinom{t-i}{j-i} \left(\tbinom{t}{i} - \nu_i \right) \, .$$

Recall also that

$$\sum_{j=1}^{t-2} (-1)^j \nu_j = 0 \, .$$

2. Decompositions of topes with respect to symmetric cycles in the tope graph, and Dehn–Sommerville type relations

Let $\mathcal{M} := (E_t, \mathcal{L}) = (E_t, \mathcal{T})$ be an arbitrary *simple* oriented matroid of rank $r \geq 3$, and let $\boldsymbol{R}$ be a *symmetric* $2t$*-cycle* in the tope graph $\mathcal{T}(\mathcal{L}(\mathcal{M}))$ of $\mathcal{M}$.

Given a tope $T \in \mathcal{T}$ of $\mathcal{M}$, consider the corresponding *decomposition*

$$T = \sum_{Q \in \boldsymbol{Q}(T,\boldsymbol{R})} Q$$

of T *with respect to the cycle* $\boldsymbol{R}$, for a *unique inclusion-minimal subset* $\boldsymbol{Q}(T, \boldsymbol{R})$ of the vertex set $\mathrm{V}(\boldsymbol{R})$ of the cycle $\boldsymbol{R}$. Suppose that

$$|\boldsymbol{Q}(T, \boldsymbol{R})| \geq 5 \ ,$$

and consider the *totally cyclic* oriented matroid

$$\mathcal{Y}_{T,\boldsymbol{R}} := {}_{-T^-}(\mathcal{M}_{\boldsymbol{R}})$$

obtained from the rank 2 oriented matroid $\mathcal{M}_{\boldsymbol{R}} := (E_t, \mathrm{V}(\boldsymbol{R}))$ with the set of topes $\mathrm{V}(\boldsymbol{R})$ by *reorientation* of $\mathcal{M}_{\boldsymbol{R}}$ on the *negative part* T^- of the tope T.

Associate with the abstract simplicial *complex*

$$\Delta := \Delta_{\mathrm{acyclic}}(\mathcal{Y}_{T,\boldsymbol{R}}) \tag{2.1}$$

of *acyclic subsets* of the ground set of the oriented matroid $\mathcal{Y}_{T,\boldsymbol{R}}$ its "*long*" f*-vector*

$$\boldsymbol{f}(\Delta; t) := \big(f_0(\Delta; t), f_1(\Delta; t), \ldots, f_t(\Delta; t)\big) \in \mathbb{N}^{t+1}$$

defined by

$$f_j(\Delta; t) := \#\{F \in \Delta \colon |F| = j\} \ , \quad 0 \leq j \leq t \ .$$

In view of (1.5), we have

$$\begin{cases} f_j(\Delta; t) = \binom{t}{j}, \ \text{if } 0 \leq j \leq 2 \ , \\ f_{t-1}(\Delta; t) = f_t(\Delta; t) = 0 \ , \\ \sum_{j=3}^{t} \left(\binom{t}{j} - f_j(\Delta; t)\right) (\mathrm{x}-1)^{t-j} = -\sum_{j=3}^{t} (-1)^j \left(\binom{t}{j} - f_j(\Delta; t)\right) \mathrm{x}^{t-j} \ . \end{cases}$$

Let us return to the symmetric cycle $\boldsymbol{R}$, and associate with the tope T the abstract simplicial complex $\boldsymbol{\Lambda} := \boldsymbol{\Lambda}(T, \boldsymbol{R})$ with the facet family

$$\{E_t - \mathbf{S}(T, Q) \colon Q \in \boldsymbol{Q}(T, \boldsymbol{R})\} \ , \tag{2.2}$$

where $\mathbf{S}(T, Q) := \{e \in E_t \colon T(e) \neq Q(e)\}$ is the *separation set* of the topes T and Q. By construction, the complex $\boldsymbol{\Lambda}$ and the complex Δ defined by (2.1) coincide, and we come to the following result:

Proposition 2.1. *Let* $\boldsymbol{R}$ *be a symmetric cycle in the tope graph of a simple oriented matroid* $\mathcal{M} := (E_t, \mathcal{T})$.

Let $T \in \mathcal{T}$ be a tope of $\mathcal{M}$ such that for the unique inclusion-minimal subset of topes $\boldsymbol{Q}(T, \boldsymbol{R}) \subset \mathrm{V}(\boldsymbol{R})$ with the property

$$\sum_{Q \in \boldsymbol{Q}(T,\boldsymbol{R})} Q = T$$

we have

$$|\boldsymbol{Q}(T, \boldsymbol{R})| \geq 5 \ .$$

(i) *The components of the long f-vector $\boldsymbol{f}(\boldsymbol{\Lambda}; t)$ of the complex*

$$\begin{aligned} \boldsymbol{\Lambda} &:= \boldsymbol{\Lambda}(T, \boldsymbol{R}) \\ &= \boldsymbol{\Lambda}(-T, \boldsymbol{R}) \end{aligned}$$

whose family of facets is defined by (2.2) *satisfy the* Dehn–Sommerville *type* relations

$$\begin{cases} f_j(\boldsymbol{\Lambda}; t) = \binom{t}{j}, \text{ if } 0 \leq j \leq 2 \ , \\ f_{t-1}(\boldsymbol{\Lambda}; t) = f_t(\boldsymbol{\Lambda}; t) = 0 \ , \\ \sum_{j=3}^{t} \left(\binom{t}{j} - f_j(\boldsymbol{\Lambda}; t) \right) (\mathrm{x}-1)^{t-j} = - \sum_{j=3}^{t} (-1)^j \left(\binom{t}{j} - f_j(\boldsymbol{\Lambda}; t) \right) \mathrm{x}^{t-j} \ . \end{cases}$$

In particular,

- *if $t = 5$, then*

$$f_3(\boldsymbol{\Lambda}; t) = \binom{t}{2} - t = 5 \ ;$$

- *if $t = 6$, then*

$$\begin{aligned} f_3(\boldsymbol{\Lambda}; t) &= \binom{t}{3} - 2t + 4 = 12 \ , \\ f_4(\boldsymbol{\Lambda}; t) &= \binom{t}{2} - 3t + 6 = 3 \ ; \end{aligned}$$

- *if $t = 7$, then*

$$\begin{array}{rclcl} f_3(\boldsymbol{\Lambda}; t) & = & \frac{1}{2} f_4(\boldsymbol{\Lambda}; t) + \binom{t}{4} - \frac{1}{2}\binom{t}{3} & = & \frac{1}{2}\big(f_4(\boldsymbol{\Lambda}; t) + 35\big) \\ & = & f_5(\boldsymbol{\Lambda}; t) + \binom{t}{4} - \binom{t}{2} + t & = & f_5(\boldsymbol{\Lambda}; t) + 21 \ , \\ f_4(\boldsymbol{\Lambda}; t) & = & 2 f_3(\boldsymbol{\Lambda}; t) - 2\binom{t}{4} + \binom{t}{3} & = & 2 f_3(\boldsymbol{\Lambda}; t) - 35 \\ & = & 2 f_5(\boldsymbol{\Lambda}; t) + \binom{t}{3} - 2\binom{t}{2} + 2t & = & 2 f_5(\boldsymbol{\Lambda}; t) + 7 \ , \\ f_5(\boldsymbol{\Lambda}; t) & = & f_3(\boldsymbol{\Lambda}; t) - \binom{t}{4} + \binom{t}{2} - t & = & f_3(\boldsymbol{\Lambda}; t) - 21 \\ & = & \frac{1}{2} f_4(\boldsymbol{\Lambda}; t) - \frac{1}{2}\binom{t}{3} + \binom{t}{2} - t & = & \frac{1}{2}\big(f_4(\boldsymbol{\Lambda}; t) - 35\big) + 14 \ ; \end{array}$$

the quantity $f_4(\boldsymbol{\Lambda}; 7)$ is odd.

(ii) *For any j, $3 \leq j \leq t-2$, we have*

$$\binom{t}{j} - f_j(\boldsymbol{\Lambda}; t) = - \sum_{i=3}^{j} (-1)^i \binom{t-i}{j-i} \left(\binom{t}{i} - f_i(\boldsymbol{\Lambda}; t) \right) \ .$$

(iii) *We have*

$$\sum_{j=1}^{t-2} (-1)^j f_j(\boldsymbol{\Lambda}; t) = 0 \ .$$

References

[1] *Björner A.*, *Las Vergnas M.*, *Sturmfels B.*, *White N.*, *Ziegler G.M.* Oriented matroids. Second edition. Encyclopedia of Mathematics, 46. – Cambridge: Cambridge University Press, 1999.

[2] *Gainanov D.N.* Graphs for pattern recognition. Infeasible systems of linear inequalities. – Berlin: De Gruyter, 2016.

[3] *Matveev A.O.* Pattern recognition on oriented matroids. – Berlin: De Gruyter, 2017.

E-mail address: andrey.o.matveev@gmail.com